



A note about conditional Ornstein-Uhlenbeck processes

Amel Bentata

► To cite this version:

| Amel Bentata. A note about conditional Ornstein-Uhlenbeck processes. 2008. hal-00211782

HAL Id: hal-00211782

<https://hal.science/hal-00211782>

Preprint submitted on 21 Jan 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A note about conditional Ornstein-Uhlenbeck processes

Amel Bentata*

11 janvier 2008

Abstract

In this short note, the identity in law, which was obtained by P. Salminen [6], between on one hand, the Ornstein-Uhlenbeck process with parameter γ , killed when it reaches 0, and on the other hand, the 3-dimensional radial Ornstein-Uhlenbeck process killed exponentially at rate γ and conditioned to hit 0, is derived from a simple absolute continuity relationship.

Keywords: Ornstein-Uhlenbeck process, Doob's h-transform, absolute continuity relationship.

Mathematical Subject Classification: 60J60; 60G15.

e-mail address: bentata@clipper.ens.fr

- All probability distributions considered in this note are defined on $\mathcal{C}(\mathbb{R}_+, \mathbb{R})$, where $(X_t, t \geq 0)$ denotes the coordinate process, and $\mathcal{F}_t = \sigma\{X_s, s \leq t\}$, its natural filtration.
- For $\gamma > 0$, and $a > 0$, we denote by \mathbb{P}_a^γ the law of the Ornstein-Uhlenbeck process with parameter γ , starting from a , i.e : under \mathbb{P}_a^γ , one has :

$$X_t = a + B_t - \gamma \int_0^t X_s ds \quad (1)$$

for a Brownian motion $(B_t, t \geq 0)$, starting from 0.

*Université Paris 6, Laboratoire de Probabilités et Modèles Aléatoires, CNRS-UMR 7599, 16, rue Clisson, 75013 Paris Cedex, France

Note that $(X_t \exp(\gamma t), t \geq 0)$ is a \mathbb{P}_a^γ martingale which is indeed equal to :

$$a + \int_0^t \exp(\gamma s) dB_s, \quad t \geq 0, \quad (2)$$

thus leading us to Doob's well-known representation of the Ornstein-Uhlenbeck process (see [2]) :

$$X_t = e^{-\gamma t} \left(a + \beta \left(\frac{e^{2\gamma t} - 1}{2\gamma} \right) \right), \quad t \geq 0, \quad (3)$$

for another Brownian motion $(\beta(u), u \geq 0)$, starting from 0.

- In this note, the 3-dimensional Ornstein-Uhlenbeck process (\vec{X}_t) with parameter γ and starting from \vec{a} , i.e : a solution of (1) where (B_t) is replaced by a 3-dimensional Brownian motion (\vec{B}_t) , and its radial part $R_t = |\vec{X}_t|$ play an important role; (R_t) solves the *SDE* :

$$R_t = a + \tilde{B}_t + \int_0^t \frac{ds}{R_s} - \gamma \int_0^t R_s ds \quad (4)$$

where $a = |\vec{a}|$ and $(\tilde{B}_t, t \geq 0)$ is a 1-dimensional Brownian motion, starting from 0.

- The main result of this note is the following :

Proposition 1. *Define the probability \mathbb{Q}_a^γ via :*

$$\mathbb{Q}_a^\gamma|_{\mathcal{F}_t} = \frac{X_{t \wedge T_0}}{a} e^{\gamma t} \cdot \mathbb{P}_a^\gamma|_{\mathcal{F}_t} \quad (5)$$

Then, \mathbb{Q}_a^γ is the law of the 3-dimensional radial Ornstein-Uhlenbeck process starting from a .

Proof. Due to Girsanov's theorem, and (2), (B_t) considered under \mathbb{Q}_a^γ solves :

$$B_t = \tilde{B}_t + \int_0^{t \wedge T_0} \frac{\exp(\gamma s)}{X_s e^{\gamma s}} ds \quad (6)$$

Now, noting that $T_0 = \infty$, \mathbb{Q}_a^γ a.s, we obtain that X_t under \mathbb{Q}_a^γ satisfies (4), as a consequence of (2) and (6). \square

- We now make a few comments :

1. Note that, for $\gamma = 0$, (5) is nothing else but Doob's h-transform relationship between Brownian motion killed when it reaches 0 and the 3-dimensional Bessel process.
2. We now write (5) in the equivalent form :

$$1_{\{t < T_0\}} \cdot \mathbb{P}_a^\gamma |_{\mathcal{F}_t} = \frac{a}{X_t} e^{-\gamma t} \cdot \mathbb{Q}_a^\gamma |_{\mathcal{F}_t} \quad (7)$$

which is nothing else but the identity in law obtained by P. Salminen ([6], Theorem 1,(i)) between, on the left hand-side, the Ornstein-Uhlenbeck process with parameter γ , killed when it reaches 0, and on the right hand-side the 3-dimensional radial Ornstein-Uhlenbeck process killed exponentially at rate γ , and conditioned to hit 0, that is the density $(\frac{1}{X_t})$ corresponding to such a conditioning may be seen a posteriori from the formula :

$$\mathbb{Q}_a^\gamma(F_t \frac{1}{X_t}) = \mathbb{Q}_a^\gamma(\frac{1}{X_t}) \mathbb{P}_a^\gamma(F_t | t < T_0), \forall F_t \in \mathcal{F}_t, \quad (8)$$

which is a consequence of (7)

It also follows from (7) that $(\frac{1}{X_t} \exp(-\gamma t), t \geq 0)$ is a strictly local martingale with respect to \mathbb{Q}_a^γ , thus extending very simply the well-known result for the case $\gamma = 0$, when (X_t) is a 3-dimensional Bessel process.

3. Note that (7) leads us to the computation of the semi-group of the killed Ornstein-Uhlenbeck, via :

$$\mathbb{E}_a^\gamma[(\frac{e^{\gamma t}}{a}) f(X_t) 1_{\{t \wedge T_0\}}] = \mathbb{Q}_a^\gamma[f(X_t) \frac{1}{X_t}], \quad (9)$$

allowing us to recover the result in [7], p122, formula (53).

4. What about $\gamma < 0$? Formula (5) is still valid thanks to the same arguments; it is closely related to Theorem 1,(ii) of [6].
5. Related informations about (radial) Ornstein-Uhlenbeck processes, and their hitting times, may be found in [1], [3], [4], [5].

Acknowledgments : I am very grateful to P. Salminen for a number of discussions during the preparation of this note.

References

- [1] L. ALILI, P. PATIE and J.L PEDERSEN (2005). Representations of the first hitting time density of an Ornstein-Uhlenbeck process. *Stoch.Models.* **21** , no. 4, 967-980.
- [2] J.L. DOOB (April 1942). The Brownian Movement and Stochastic Equations. *The Annals of Mathematics.* **43** , no. 2, 351-369.
- [3] K.D. ELWORTHY , X.-M LI and M. YOR (1999). The importance of strictly local martingales; applications to radial Ornstein-Uhlenbeck processes. *Probability theory and related fields.* **115** , no. 3, 325-355.
- [4] A. GOING-JAESCHKE and M. YOR (April 2003). A clarification note about hitting times densities for Ornstein-Uhlenbeck processes. *Finance and Stochastics.* **91** , no. 1.
- [5] A. GOING-JAESCHKE and M. YOR (2003). A survey and some generalizations of Bessel processes. *Bernoulli.* **9**, 313-349.
- [6] P. SALMINEN (December 1984). On conditional Ornstein-Uhlenbeck Processes. *Advances in Applied Probability.* **16** , no. 4.
- [7] P. SALMINEN , P. VALLOIS and M. YOR (December 2006). On the excursion theory for linear diffusions. *Japanese Journal of Mathematics.* **91** , no. 1, 97-127.